

OPTICAL SPECTROPOLARIMETRY OF SN 2002AP: A HIGH VELOCITY ASYMMETRIC EXPLOSION¹

K. S. KAWABATA^{2,3}, D. J. JEFFERY⁴, M. IYE^{2,5}, Y. OHYAMA⁶, G. KOSUGI⁶, N. KASHIKAWA²,
 N. EBIZUKA⁷, T. SASAKI⁶, K. SEKIGUCHI⁶, K. NOMOTO^{8,9}, P. MAZZALI^{9,8,10}, J. DENG^{9,8},
 K. MAEDA⁸, K. AOKI⁶, Y. SAITO², T. TAKATA⁶, M. YOSHIDA¹¹, R. ASAI⁸, M. INATA¹¹,
 K. OKITA¹¹, K. OTA^{8,2}, T. OZAWA¹², Y. SHIMIZU¹¹, H. TAGUCHI¹³, Y. YADOMARU¹²,
 T. MISAWA^{8,2}, F. NAKATA^{8,2}, T. YAMADA², I. TANAKA², AND T. KODAMA¹⁴
accepted for publication in the Astrophysical Journal (Letters)

ABSTRACT

We present spectropolarimetry of the Type Ic supernova SN 2002ap and give a preliminary analysis: the data were taken at two epochs, close to and one month later than the visual maximum (2002 February 8). In addition we present June 9 spectropolarimetry without analysis. The data show the development of linear polarization. Distinct polarization profiles were seen only in the O I $\lambda 7773$ multiplet/Ca II IR triplet absorption trough at maximum light and in the O I $\lambda 7773$ multiplet and Ca II IR triplet absorption troughs a month later, with the latter showing a peak polarization as high as $\sim 2\%$. The intrinsic polarization shows three clear position angles: 80° for the February continuum, 120° for the February line feature, and 150° for the March data. We conclude that there are multiple asymmetric components in the ejecta. We suggest that the supernova has a bulk asymmetry with an axial ratio projected on the sky that is different from 1 by order 10%. Furthermore, we suggest very speculatively that a high velocity ejecta component moving faster than $\sim 0.115c$ (e.g., a jet) contributes to polarization in the February epoch.

Subject headings: polarization — supernovae: individual (SN 2002ap)

1. INTRODUCTION

SN 2002ap was discovered in the nearby spiral galaxy M74 (= NGC 628) on 29 January 2002 (Nakano et al. 2002) and reached its maximum of $V \sim 12.4$ mag on February 8 (Gal-Yam, Ofek, & Shemmer 2002). It has been classified as a Type Ic supernova (SN Ic) and suggested to be a hypernova (but at the low-energy end of the sequence of hypernovae; Mazzali et al. 2002 and references therein).

A SN Ic is thought to be the result of the core collapse of a massive star that has either lost its hydrogen and helium envelopes prior to the explosion or has an invisible helium envelope due to low excitation. The details of the explosion mechanism are still under discussion (Nomoto, Iwamoto, & Suzuki 1995; Branch 2001 and references therein). SN 1998bw, the most luminous and energetic Type Ic ‘hypernova’ to date, has been particularly well studied, and its probable connection with the γ -ray burst GRB 980425 has been pointed out (e.g., Galama et al. 1998; Iwamoto et al. 1998; Nomoto et al. 2001). An

aspherical hyperenergetic explosion has been suggested to explain the slowly-declining light curve of SN 1998bw and the narrowness of the [O I] $\lambda 6300, 6363$ emission line in the nebular phase (Mazzali et al. 2001; Nakamura et al. 2001; Maeda et al. 2002).

Intrinsic polarization is zero for SNe (which are unresolved sources) if they are spherically symmetric: any intrinsic polarization thus reveals asymmetry. Core-collapse supernovae are, in fact, generally polarized in the continuum at levels of $p \simeq 0.5\text{--}4\%$ due to electron scattering and their polarization increases after optical maximum light (e.g., Jeffery 1991b; Wang et al. 1996, 2001; Leonard et al. 2001); however, the polarization falls to zero at very late times, when the electron scattering opacity becomes very low (e.g., Jeffery 1991b). The typical line polarization profile—actually mostly due to electron polarized light interacting with a line—predicted theoretically (Jeffery 1989) and to some degree confirmed observationally in SN 1987A and other supernovae (Jeffery 1991a,b; Leonard

¹ Based on data obtained at the Subaru Telescope, which is operated by the National Astronomical Observatory of Japan (NAOJ)

² Opt. & IR Astron. Div., NAOJ, Mitaka, Tokyo 181-8588, Japan

³ E-mail: koji.kawabata@nao.ac.jp

⁴ Dep. of Phys., New Mexico Institute of Mining and Technology, Socorro, NM 87801, USA

⁵ Dep. of Astron., Graduate Univ. for Advanced Studies, Mitaka, Tokyo 181-8588, Japan

⁶ Subaru Telescope, NAOJ, 650 North A’ohoku Place, Hilo, HI 96720, USA

⁷ RIKEN, Wako, Saitama 351-0198, Japan

⁸ Dep. of Astron., Univ. of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

⁹ Research Center for the Early Universe, Univ. of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

¹⁰ Osservatorio Astronomico, Via Tiepolo, 11, 34131 Trieste, Italy

¹¹ Okayama Astrophys. Obs., NAOJ, Asakuchi-gun, Okayama 719-0232, Japan

¹² Misato Obs., Amakusa-gun, Wakayama 640-1366, Japan

¹³ Dep. of Astron. & Earth Sci., Tokyo Gakugei University, Koganei, Tokyo 184-8501, Japan

¹⁴ Theory Div., NAOJ, Mitaka, Tokyo 181-8588, Japan

& Filippenko 2001; Leonard et al. 2001), is an inverted P Cygni profile: strong polarization maximum at the flux P Cygni trough feature and polarization minimum at the flux P Cygni emission feature. Line blending and other intrinsic effects can distort these profiles. For SN 1998bw, an intrinsic optical polarization of 0.4–0.6 % was found, suggesting an asymmetry of less than 2/1 in the axial ratio of the ejecta (Patat et al. 2001; Kay et al. 1998). No distinct line polarization features were seen probably due to the poor S/N or the relatively narrow wavelength range of those observations.

2. OBSERVATIONS AND DATA REDUCTION

The spectropolarimetry was taken with the 8.2-m Subaru Telescope equipped with the Faint Object Camera and Spectrograph (FOCAS, Kashikawa et al. 2000; Yoshida et al. 2000) from 2002 February 9 through June 9. For February and June observations, we used a 300 grooves mm^{-1} grism (the central wavelength of 5500 Å) with a $0''.4$ width slit, resulting a spectral resolution ($\lambda/\Delta\lambda$) was ~ 1200 . For March observation, we used another 300 gr mm^{-1} grism (7500 Å) with a $0''.8$ width slit, resulting in $\lambda/\Delta\lambda \sim 650$. The nightly total exposure time was 960 to 4000 sec. The linear polarimetric module of FOCAS consists of a rotating superachromatic half-wave plate and a crystal quartz Wollaston prism, and both the ordinary and the extraordinary rays are simultaneously recorded on two MIT/LL CCDs ($2k \times 4k \times 15\mu\text{m}$). A typical observing sequence consisted of four integrations at the $\psi = 0^\circ, 45^\circ, 22.5^\circ$ and 67.5° positions of the half-wave plate. Stokes Q/I and U/I were calculated as in §6.1.2 of Tinbergen (1996). For polarimetric calibration, we obtained data for unpolarized and polarized standard stars, including measurements of flatfield lamps through fully-polarizing filters. The flux was calibrated using observations of G191B2B and BD+28°4211 (Oke 1990), and then was multiplied by a constant to match the VSNET¹⁴ photometric data.

3. RESULTS AND DISCUSSION

Figure 1 shows the observed flux and polarization spectra. Several blueshifted broad absorption lines can be identified in the February flux spectrum (Mazzali et al. 2002). For March and June flux spectra a detailed analysis has yet to be done. However, we note that the March spectra resemble that of SN 1997ef at day 67 (Mazzali, Iwamoto, & Nomoto 2000), including the onset of net emission in Ca II $\lambda\lambda 8498, 8542, 8662$ (i.e., the Ca II IR triplet). The significant emission line at $\sim 6300\text{Å}$ in the June spectrum is identified as [O I] $\lambda 6300, 6363$ as in SN 1997ef (Mazzali et al. 2001). The polarization is $\gtrsim 0.5\%$ at a position angle (PA) of $\theta = 120^\circ \pm 20^\circ$ over the observed wavelengths. Significant day-by-day variation is not seen in the polarization spectra within each month. Here we will only analyze monthly averages.

3.1. Interstellar and Intrinsic Polarization

Interstellar polarization (ISP) varies slowly with wavelength in the optical and is well approximated by the empirical formula $p_{\text{ISP}}(\lambda) = p_{\text{max}} \cdot \exp[-1.15 \ln^2(\lambda_{\text{max}}/\lambda)]$,

where $p_{\text{max}} = p_{\text{ISP}}(\lambda_{\text{max}})$ is the peak polarization (Serkowski, Mathewson, & Ford (1975), hereafter SMF). In March the emission feature of the Ca II IR triplet line profile shows strong net emission due to NLTE processes. Such NLTE line flux is necessarily unpolarized on emission. Since the line profile is still broad (absorption minimum at a $\sim -14,000 \text{ km s}^{-1}$ redshift, corresponding to an enclosed mass of $2 M_\odot$ in model CO100/4, which has a total mass of $2.4 M_\odot$ (Mazzali et al. 2002)), much of the emission is probably coming from far out in the ejecta, where the electron optical depth is low. We conclude that the flux from this emission line is mostly unscattered by electrons and unpolarized, and that it dilutes the polarized electron scattered flux. If this were the only effect, then the intrinsic polarization should show a distinct minimum nearly exactly at the wavelength of the flux emission maximum. Since, in fact, the polarization is roughly constant across the P Cygni emission feature (8400–9000 Å), apart from small variations that may be mostly noise, the line is not only diluting the polarized flux but is also probably strongly depolarizing it. The intrinsic polarization across the emission feature is probably close to zero. We will assume that the observed polarization of the emission feature is the ISP: thus $p_{\text{ISP}}(8600 \text{ Å}) \approx 0.5\%$. We next adopt the median $\lambda_{\text{max}} = 5370 \text{ Å}$ found by SMF for 30 stars with $p_{\text{max}}/E_{B-V} \geq 7.0$. Then from a non-linear regression we find $p_{\text{max}} = 0.64 \pm 0.20\%$ and $\theta_{\text{ISP}} = 120^\circ \pm 10^\circ$. (The uncertainties are crude estimates based on the alternative assumption that only line flux dilution, and not line depolarization, occurs in the region of the Ca II IR triplet emission feature.) Since Takada-Hidai, Aoki, & Zhao (2002) derive a color excess for SN 2002ap of $E_{B-V} = 0.09$ (a sum of 0.07 within our Galaxy and 0.02 within M74) from interstellar Na D absorptions, our assumption of the SMF λ_{max} is consistent: $p_{\text{max}}/E_{B-V} = 0.64/0.09 \approx 7$.

The estimated ISP (EISP) is consistent with other factors. In an ISP catalog (Heiles 2000), 16 stars are within 10° of SN 2002ap. The data for these stars suggest a possible positive correlation between polarization and the distance along the line of sight toward SN 2002ap. The two most distant stars among them, HD8919 ($d = 525 \text{ pc}$) and HD9560 ($d = 437 \text{ pc}$) show $(p, \theta) = (0.32 \pm 0.10\%, 99^\circ \pm 9^\circ)$ and $(0.48 \pm 0.09\%, 123^\circ \pm 5^\circ)$, respectively. On the other hand, it has been found that $p_{\text{max}}(\%)$ has an empirical upper limit of $9E_{B-V}$ (SMF). From the derived $E_{B-V} = 0.09$, an upper limit on the ISP toward the supernova is 0.81 %. The EISP is nicely sandwiched between the possible lower bounds of the cited stars and the empirical upper limit. The EISP position angle is also consistent with the position angle of the spiral arm in M74 at the position of SN 2002ap, $110^\circ\text{--}140^\circ$ (see DPOSS images). The June 9 polarization spectrum has polarization consistent with the EISP at the peak of the strong emission flux of the [O I] $\lambda 6300, 6363$ forbidden line (Figure 1c): one would expect that polarization to be ISP for a strong emission line in the nebular epoch. (The June 9 data has low S/N: in the following, we analyze only the higher S/N polarization spectra of the earlier two epochs.)

Figure 2 shows the intrinsic (i.e., EISP-subtracted) polarization plotted on a QU diagram: the polarization

¹⁴ <http://www/kusastro.kyoto-u.ac.jp/vsnet>

points are connected according to their wavelength ordering. Given the uncertainty in the EISP, points within 0.2 % of the origin must be considered very uncertain.

If one assumes that the intrinsic supernova polarization is produced by a single axisymmetric component in the ejecta, then the intrinsic polarization plotted on a QU diagram should lie on a line passing through the origin. It can be seen that the polarization in February has two clear position angles, PA less than or $\sim 120^\circ$ (associated with the O I/Ca II line trough) and $PA \sim 80^\circ \pm 20^\circ$ (associated with the continuum from $\sim 5700\text{--}8200\text{ \AA}$), joined by a somewhat complicated transition. The polarization in March has a clear position angle $PA \sim 150^\circ$ associated with the Ca II line trough and with at least some of the continuum. We conclude that there are multiple asymmetric components, and that their contribution varies with time. It is likely that the recession of the supernova photosphere uncovers different asymmetries.

3.2. Possible Models

Figures 3a, b, and c show the flux and intrinsic polarization corrected for heliocentric redshift [$v_{\text{helio}} = +631\text{ km s}^{-1}$ (Smartt & Meikle 2002)] and interstellar extinction ($E_{B-V} = 0.09$). The figures show that polarization is low and barely significant (given the uncertainty in the EISP), except in the regions $\sim 6700\text{--}8000\text{ \AA}$ for February and $\sim 6700\text{--}8300\text{ \AA}$ for March.

The low continuum polarization blueward of $\sim 6700\text{ \AA}$ in both epochs may be due to the depolarizing effect of lines (Howell et al. 2001): in supernova spectra, lines generally become stronger further to the blue. The continuum polarization, where it is significant (a relatively small region) is $\sim 0.4\%$ in both epochs. If the asymmetry is assumed to be an axisymmetric, global prolate or oblate asymmetry, then the continuum polarization can be explained by an axial ratio (assuming there is a main axis) projected on the sky that is different from 1 by of order 10 %. This estimate is a crude one based on realistic, but parameterized, calculations (Höflich 1991, 1995). The estimate is also crude because, as noted above, the asymmetry cannot be completely axisymmetric. The estimated asymmetry is not large, compared to those estimated for some other supernovae (e.g., Wang et al. 2001).

The three distinct line polarization profiles seen in Figure 3 (at the O I/Ca II flux absorption in February and the O I and Ca II flux absorptions in March) can partially be accounted for by the inverted P Cygni profile (see § 1): the polarization maxima associated with line trough features are clear. Without detailed modeling more information probably cannot be extracted from these profiles.

For the February continuum polarization, we suggest a radically different origin from the bulk asymmetry assumed in prolate/oblate models or element inhomogeneity models (see below). In Figure 3d we show the intrinsic polarized flux ($p \times F$) compared to the observed flux scaled down by a factor of 0.0018 and non-relativistically redshifted by a velocity of $0.23c$: i.e., $\lambda_{\text{redshifted}} = \lambda/0.77$. There is fair agreement over the range $\sim 5000\text{--}8000\text{ \AA}$. In response to this comparison, Leonard et al. (2002) made a similar comparison for their SN 2002ap spectropolarimetry and also obtained a similarly good agreement. The agreement suggests (but does not prove) that a large com-

ponent of polarized flux comes from electron scattering in an ionized clump (i.e., a jet) thrown out of the supernova explosion. (We assume a single jet or clump for simplicity here, although a pair of bipolar jets are a physical possibility (e.g., Wheeler, Meier, & Wilson 2002).) In a simple non-relativistic picture, the scattered light is redshifted by $v_{\text{red}} \sim v_{\text{jet}}(1 + \cos i)$, where, $v_{\text{red}} = 0.23c$, v_{jet} is the characteristic velocity of scattering relative to the supernova center, and i is the inclination angle of jet to the line of sight measured from the far side of the supernova. The jet polarization component is calculated from

$$p_{\text{jet}}(\lambda) = f \cdot \frac{F[\lambda(1 - v_{\text{red}}/c)]}{F(\lambda)},$$

where $F(\lambda)$ is the corrected flux, $1 - v_{\text{red}}/c$ is the blueshift back to the origin of the scattered flux observed at λ , and the scale factor $f = 0.0018$: f accounts both for the polarization of the jet scattered flux and the fraction scattered. The polarization is wavelength dependent even though electron scattering is wavelength-independent since the scattered flux comes from a bluer part of the spectrum than the part it is added to. Electron scattering depends on scattering direction: e.g., maximum polarization occurs for $i = 90^\circ$; half as much for $i = 45^\circ$ or 135° ; zero for $i = 0^\circ$ or 180° . The jet velocities corresponding to $i = 90^\circ$, 45° , and 0° are $0.23c$, $0.135c$, and $0.115c$, respectively. Thus $0.115c$ is a lower bound on the jet velocity and $i \gtrsim 90^\circ$ would require a somewhat relativistic jet.

High velocity jet-like clumps have been proposed in some hydrodynamic explosion models for SNe and GRB's (e.g., Nagataki et al. 1997; MacFadyen & Woosley 1999; Maeda et al. 2002; Wheeler, Meier, & Wilson 2002). If a jet is thrown out of an exploding supernova core, then it is plausible that it carries some radioactive ^{56}Ni . The gamma-rays from decay would keep the jet ionized to some degree just as they keep the nebular phase bulk ejecta ionized.

If the jet picture is correct, then the position angle of the jet on the sky is $\sim 170^\circ$ (or $\sim 350^\circ$) since the jet polarization component has position angle $\sim 80^\circ$ and electron scattering polarizes perpendicularly to the scattering plane. The O I/Ca II line polarization maximum in the February data cannot easily be associated with the jet. As Figure 2 shows, the observed position angle makes an excursion from $\sim 80^\circ$ up to 120° across the polarization maximum. Some of the line polarization may arise in the bulk asymmetry of the supernova. It is possible that the position angle of $\sim 120^\circ$ is the net result of a jet polarizing at $\sim 80^\circ$ and a bulk asymmetry polarizing at $\sim 150^\circ$ (i.e., at the position angle observed in the March data). To test this model we have eliminated the jet polarization component from the February intrinsic polarization by subtracting jet polarization Stokes parameters from the intrinsic polarization Stokes parameters. The residual polarization and position angle spectra are plotted in Figures 3e,f. The position angle of the residual polarization for February from the region of significant polarization (i.e., $\sim 6700\text{--}8000\text{ \AA}$) is now approximately centered on 150° and deviates by more than 30° only in a few isolated points. The results in Figure 3f are thus consistent with the jet model.

The jet may not be a completely separated amount of ejecta, but rather a blob rich in ^{56}Ni moving at $\gtrsim 0.115c$.

A high-velocity, ^{56}Ni -rich region is required both in the hypernova explosion models of SN 1998bw (Nakamura et al. 2001; Maeda et al. 2002), and in SN 2002ap (Mazzali et al. 2002) in order to reproduce the light curves. Spectral synthesis for SN 2002ap (Mazzali et al. 2002) suggests some material at velocities higher than the photospheric velocities of $\sim 0.1c$ – $0.117c$ (Mazzali et al. 2002; Kinugasa et al. 2002) up to $0.22c$. The ionization of the blob would likely be increased by radioactive ^{56}Ni and this would likely make the blob more polarizing than other parts of the ejecta at the same velocity leading to net polarization. Other chemical inhomogeneities in the ejecta at varying velocities are also possible (Maeda et al. 2002) and would affect polarization in complicated ways.

4. CONCLUDING REMARKS

Our results are summarized in the Abstract. More realistic modeling is necessary for a more definitive understanding of the polarization. The degree of the bulk asymmetry suggested in this paper may be tested with the line widths and their ratios in the nebular spectrum (Maeda et al. 2002).

We are grateful to the staff members at the Subaru Telescope for their kind help and their rearrangement of the telescope maintenance schedule for our observation on March 8. This work has been supported in part by the grant-in-Aid for Scientific Research (12640233, 14047206, 14540223) and COE research (07CE2002) of the Ministry of Education, Science, Culture, Sports, and Technology in Japan.

REFERENCES

- Branch, D. 2001, in *Supernovae and Gamma-Ray Bursts*, ed. M. Livio, N. Panagia and K. Sahu (Cambridge: Cambridge Univ. Press), 96
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Galama, T. J., et al. 1998, *Nature*, 395, 670
- Gal-Yam, A., Ofek, E. O., & Shemmer, O. 2002, *MNRAS*, 332, L73
- Heiles, C. 2000, *AJ*, 119, 923
- Höflich, P. 1991, *A&A*, 246, 481
- Höflich, P. 1995, *ApJ*, 440, 821
- Howell, D. A., Höflich, P., Wang, L., & Wheeler, J. C. 2001, *ApJ*, 556, 302
- Iwamoto, K., et al. 1998, *Nature*, 395, 672
- Jeffery, D. J. 1989, *ApJS*, 71, 951
- Jeffery, D. J. 1991a, *ApJ*, 375, 264
- Jeffery, D. J. 1991b, *ApJS*, 77, 405
- Kashikawa, N., et al. 2000, *Proc. SPIE*, 4008, 104
- Kay, L. E., Halpern, J. P., Leighly, K. M., Heathcote, S., & Magalhaes, A. M. 1998, *IAU Circ.*, No. 6969
- Kinugasa, K., et al. 2002, *ApJ*, 577, L97
- Leonard, D. C., Filippenko, A. V. 2001, *PASP*, 113, 920
- Leonard, D. C., Filippenko, A. V., Ardila, D. R., & Brotherton, M. S. 2001, *ApJ*, 553, 861
- Leonard, D. C., Filippenko, A. V., Chornock, R., & Foley, R. J. 2002, *PASP*, in press
- MacFadyen, A. I., & Woosley, S. E. 1999, *ApJ*, 524, 262
- Maeda, K., Nakamura, T., Nomoto, K., Mazzali, P. A., Patat, F., & Hachisu, I. 2002, *ApJ*, 565, 405
- Mazzali, P. A., Iwamoto, K., & Nomoto, K. 2000, *ApJ*, 545, 407
- Mazzali, P. A., Nomoto, K., Patat, F., & Maeda, K. 2001, *ApJ*, 559, 1047
- Mazzali, P. A., et al. 2002, *ApJ*, 572, L61
- Nagataki, S., Hashimoto, M., Sato, K., & Yamada, S. 1997, *ApJ*, 486, 1026
- Nakamura, T., Mazzali, P. A., Nomoto, K., & Iwamoto, K. 2001, *ApJ*, 550, 991
- Nakano, S., & Hirose, Y. 2002, *IAU Circ.*, No. 7810
- Nomoto, K., Iwamoto, K., & Suzuki, T. 1995, *Phys. Rep.*, 256, 173
- Nomoto, K., et al. 2001, in *Supernovae and Gamma-Ray Bursts*, ed. M. Livio, N. Panagia and K. Sahu (Cambridge: Cambridge Univ. Press), 144
- Oke, J. B. 1990, *AJ*, 99, 1621
- Patat, F., et al. 2001, *ApJ*, 555, 900
- Serkowski, K., Mathewson, D. S., & Ford, V. L. 1975, *ApJ*, 196, 261 (SMF)
- Smartt, S., & Meikle, P. 2002, *IAU Circ.*, No. 7822
- Takada-Hidai, M., Aoki, W., & Zhao, G. 2002, in preparation
- Tinbergen, J. 1996, *Astronomical Polarimetry* (New York: Cambridge Univ. Press)
- Wang, L., Wheeler, J. C., Li, Z., & Clocchiatti, A. 1996, *ApJ*, 467, 435
- Wang, L., Howell, D. A., Höflich, P., & Wheeler, J. C. 2001, *ApJ*, 550, 1030
- Wheeler, J. C., Meier, D. L., & Wilson, J. R. 2002, *ApJ*, 568, 807
- Yoshida, M., et al. 2000, *Proc. SPIE*, 4009, 240

FIG. 1.— Flux and polarization spectra of SN 2002ap. Heliocentric redshift, interstellar extinction and polarization have not been corrected for. From top to bottom, we plot (a) total flux, (b, c) polarization level p and position angle θ on each observation day as indicated. The polarimetric data are binned to a constant photon noise of 0.05 % which is shown by the error bars of polarization points. The EISP component is shown by a dashed curve in (b,c).

FIG. 2.— QU -diagram of the monthly-averaged intrinsic (i.e., ISP-subtracted) polarization for February and March epochs. The data are binned to a constant photon noise of 0.04 %. It can be seen that the polarization in February has, at least, two preferred axes: $PA \sim 120^\circ$ (associated with the O I/Ca II line trough) and $PA \sim 80^\circ$ (associated with the continuum). The polarization in March has a clear position angle $PA \sim 150^\circ$ associated with the Ca II line trough and the significantly polarized continuum. These position angles are indicated by thick arrows. Note that the position angle on the sky is half the angular location on a QU diagram.

FIG. 3.— Intrinsic polarization spectra corrected for heliocentric redshift and interstellar extinction. From top to bottom, we plot (a) total flux in $\text{erg s}^{-1} \text{cm}^{-2} \text{\AA}^{-1}$, (b) polarization level p , (c) position angle θ , (d) polarized flux, and (e,f) p and θ of the residual polarization after the jet polarization component has been subtracted from the February data. The February flux is the mean of Feb 9 and 11, and the March flux is the mean of Mar 8 and 10. We adopt the normal interstellar extinction curve (Cardelli, Clayton, & Mathis 1989). Deep absorption bands due to the terrestrial atmosphere and the interstellar medium have been removed by interpolation using nearby continuum levels. The polarimetric data are binned in the same manner as in Fig. 2. The solid curve in (d) is the February flux multiplied by 0.0018 and redshifted by +0.23c (see §3.2).





